

## Contents

### iii Graphical Abstracts

#### Regular Papers

- |  |     |  |
|--|-----|--|
| C.J. Cramer, D.G. Truhlar (Minneapolis, MN, USA) and A.D. French (New Orleans, LA, USA)  | 1   | Exo-anomeric effects on energies and geometries of different conformations of glucose and related systems in the gas phase and aqueous solution  |
| M. Martín-Pastor, J.F. Espinosa, J.L. Asensio and J. Jiménez-Barbero (Madrid, Spain)   | 15  | A comparison of the geometry and of the energy results obtained by application of different molecular mechanics force fields to methyl $\alpha$ -lactoside and the C-analogue of lactose |
| T. Yazaki, M. Ohnishi, S. Rokushika (Kyoto, Japan) and G. Okada (Shizuoka, Japan)  | 51  | Subsite structure of the $\beta$ -glucosidase from <i>Aspergillus niger</i> , evaluated by steady-state kinetics with cello-oligosaccharides as substrates                               |
| K.L. Larsen, F. Mathiesen and W. Zimmermann (Aalborg, Denmark)   | 59  | Separation and analysis of cyclodextrins by capillary zone electrophoresis   |
| T. Mori, S. Fujita and Y. Okahata (Yokohama, Japan)  | 65  | Transglycosylation in a two-phase aqueous–organic system with catalysis by a lipid-coated $\beta$ -D-galactosidase   |
| P.L. Barili, G. Catelani, F. D'Andrea, F. De Rensis and P. Falcini (Pisa, Italy)   | 75  | Improved preparation of 2,3:5,6:3',4'-tri- <i>O</i> -isopropylidenelactose dimethyl acetal and its 6'- <i>O</i> -(1-methoxy-1-methylethyl) derivative                                    |
| E. Fanton, C. Fayet and J. Gelas (Aubière, France)   | 85  | Long-chain acetals derived from sucrose as a new class of surfactants  |
| M.A.E. Sallam (Alexandria, Egypt), E.I. Ibrahim (Ismailia, Egypt), K.A.A. El-Eter (Alexandria, Egypt) and J.M. Cassady (Columbus, OH, USA) | 93  | Synthesis and anomeric configuration of 2-(erythrofuransyl)benzimidazole C-nucleoside analogues  |
| A. Radha and R. Chandrasekaran (West Lafayette, IN, USA)   | 105 | X-Ray and conformational analysis of arabinan  |
| M. Hisamatsu, T. Mishima, K. Teranishi and T. Yamada (Tsu, Japan)  | 117 | The correlation between adhesion of schizophyllan to yeast glucan and its effect on regeneration of yeast protoplast   |

#### Notes

- |   |     |   |
|---|-----|---|
| S.Y. Hassan, H.M. Faidallah, A.M. El-Massry and M.M. El-Sadek (Alexandria, Egypt)           | 123 | Synthesis and reactions of 5-(D-arabino-tetrahydroxybutyl)-3-(2,3-dihydro-1,3,4-oxadiazole-2-thion-5-yl)-2-methylfuran and 5-(D-arabino-tetrahydroxybutyl)-3-(2-substituted amino-1,3,4-oxadiazol-5-yl)-2-methylfuran |
| S. Cottaz (Grenoble, France), P. Rollin (Orléans, France) and H. Driguez (Grenoble, France) | 127 | Synthesis of 2-deoxy-2-fluoro-glucotropaeolin, a thioglucosidase inhibitor  |

### iii Graphical Abstracts

#### Regular Papers

- |   |     |   |
|---|-----|---|
| N. Wang and G.R. Gray (Minneapolis, MN, USA)  | 131 | Reductive cleavage of permethylated polysaccharides with borane–methyl sulfide complex and butyltin trichloride |
| G. Blond, D. Simatos (Dijon, France), M. Catté, C.G. Dussap and J.B. Gros (Aubière, France) | 139 | Modeling of the water–sucrose state diagram below 0 °C  |

M. Kluge (Leipzig, Germany), B. Schneider (Halle / S., Germany) and D. Sicker (Leipzig, Germany)	147	Diastereoselective synthesis of the benzoxazinone acetal glucoside <i>ent</i> -GDIMBOA: the first enantiomer of a natural acetal glucoside
S. Bay (Paris, France), O. Berthier-Vergnes (Lyon, France) and D. Cantacuzene (Paris, France)	153	Synthesis of $\alpha$ - and $\beta$ -biotinylated T-antigen
A. Blaskó, C.A. Bunton (Santa Barbara, CA, USA), S. Bunel, C. Ibarra and E. Moraga (Santiago, Chile)	163	Determination of acid dissociation constants of anomers of amino sugars by $^1\text{H}$ NMR spectroscopy
J.M. Kim and R. Roy (Ottawa, Canada)	173	New prototypical O-linked-glycopeptidomimetics corresponding to the linkage region of proteoglycans
X.-L. Sun, T. Kai, H. Takayanagi and K. Furuhashi (Tokyo, Japan)	181	Syntheses of sialic acid analogues with acylamino groups at C-4 ( <i>N</i> -acyl regioisomers of sialic acids)
G. Kogan, D. Uhrin, J.-R. Brisson and H.J. Jennings (Ottawa, Canada)	191	Structural basis of the <i>Neisseria meningitidis</i> immunotypes including the L4 and L7 immunotypes
Y. Yachida, K. Tsuchihashi and S. Gasa (Sapporo, Japan)	201	Novel di- <i>O</i> -acetylated GM3s from equine erythrocytes, one containing 4,9-di- <i>O</i> -acetyl- <i>N</i> -glycolylneuraminic acid and another containing 4- <i>O</i> -acetyl- <i>N</i> -glycolylneuraminic acid and 6- <i>O</i> -acetyl-D-galactose
A.M. Winn and S.G. Wilkinson (Hull, UK)	213	Structure of the O2 antigen of <i>Stenotrophomonas</i> ( <i>Xanthomonas</i> or <i>Pseudomonas</i> ) <i>maltophilia</i>
C. Petersson (Uppsala, Sweden), W. Jachymek (Wrocław, Poland), L. Kenne (Uppsala, Sweden), T. Niedziela and C. Lugowski (Wrocław, Poland)	219	Structural studies of the O-specific chain of <i>Hafnia alvei</i> strain PCM 1190 lipopolysaccharide
N.P. Arbatsky, A.S. Shashkov (Stockholm, Sweden; Moscow, Russian Federation), G. Widmalm (Stockholm, Sweden), Y.A. Knirel (Moscow, Russian Federation), K. Zych and Z. Sidorczyk (Łódź, Poland)	229	Structure of the O-specific polysaccharide of <i>Proteus penneri</i> strain 25 containing <i>N</i> -(L-alanyl) and multiple <i>O</i> -acetyl groups in a tetrasaccharide repeating unit
L. Kandra, G. Gyémánt, E. Farkas and A. Lipták (Debrecen, Hungary)	237	Action pattern of porcine pancreatic $\alpha$ -amylase on three different series of $\beta$ -maltooligosaccharide glycosides
	I	Corrigendum
	III	Announcement
	iii	Graphical Abstracts

## Regular papers

C.R. Rozanas and G.R. Gray (Minneapolis, MN, USA)	243	Reductive cleavage of the positional isomers of benzoylated and methylated methyl $\alpha$ -D-mannopyranoside
B. Laignel, C. Bliard, G. Massiot and J.M. Nuzillard (Reims, France)	251	Proton NMR spectroscopy assignment of D-glucose residues in highly acetylated starch
Y. Kobayashi and T. Tsuchiya (Kawasaki, Japan)	261	Synthesis of 2''-oxidized derivatives of 5-deoxy-5-epi-5-fluoro-dibekacin and -arbakacin, and study on structure-chemical shift relationships of urethane(or amide)-type NH protons in synthetic intermediates
Y.M. Ahn and G.R. Gray (Minneapolis, MN, USA)	279	Diastereoselectivity in the transglycosidation of methyl 2-deoxy-3,4,6-tri- <i>O</i> -methyl-2-( <i>N</i> -methylacetamido)-D-glucopyranoside, -galactopyranoside, and -mannopyranoside with racemic 2-butanol under reductive-cleavage conditions
R. Hoos, A. Vasella (Zurich, Switzerland), K. Rupitz and S.G. Withers (Vancouver, Canada)	291	D-Glyconhydroximolactams strongly inhibit $\alpha$ -glycosidases
J.-P. Vincken, G. Beldman and A.G.J. Voragen (Wageningen, The Netherlands)	299	Substrate specificity of endoglucanases: what determines xyloglucanase activity?
J.O. Previato (Rio de Janeiro, Brasil), C. Jones (Potters Bar, UK), M.P. Stephan, L.P.A. Almeida and L. Mendonça-Previato (Rio de Janeiro, Brasil)	311	Structure of the repeating oligosaccharide from the lipopolysaccharide of the nitrogen-fixing bacterium <i>Acetobacter diazotrophicus</i> strain PAL 5

- V. Planchot, P. Colonna and A. Buleon (Nantes, France)** 319 Enzymatic hydrolysis of  $\alpha$ -glucan crystallites
- G.K. Moates, T.R. Noel, R. Parker and S.G. Ring (Norwich, UK)** 327 The effect of chain length and solvent interactions on the dissolution of the B-type crystalline polymorph of amylose in water

## Notes

- E.A. Yates, F. Santini, A. Bisio and C. Cosentino (Milan, Italy)** 335 Evidence for a heparin derivative containing an *N*-sulfated aziridine ring that retains high anti-factor Xa activity
- Z. Smiatacz, I. Chrzczanowicz and H. Myszka (Gdańsk, Poland)** 341 Chemical modification of some methyl (3,4-di-*O*-acetyl-2-deoxy-2-hydroxyimino- $\alpha$ -D-arabino-hexopyranosid)uronates

- I Erratum
- III Full Instructions to Authors
- XV Author index
- XVII Subject index
- XXI Contents

1  
3